

Problemen

| Problem Section

This Problem Section is open to everyone; everybody is encouraged to send in solutions and propose problems. Group contributions are welcome.

For each problem, the most elegant correct solution will be rewarded with a book token worth € 20. At times there will be a Star Problem, to which the proposer does not know any solution. For the first correct solution sent in within one year there is a prize of € 100. When proposing a problem, please either include a complete solution or indicate that it is intended as a Star Problem.

Please send your submission by e-mail (LaTeX is preferred), including your name and address to problems@nieuwarchief.nl.

The deadline for solutions to the problems in this edition is 1 March 2017.

Problem A (folklore)

For a finite sequence $s = (s_1, \dots, s_n)$ of positive integers, denote by $p(s)$ the number of ways to write s as a sum $s = \sum_{i=1}^n a_i e_i + \sum_{j=1}^{n-1} b_j (e_j + e_{j+1})$ with all a_i and b_j non-negative. Here e_i denotes the sequence of which the i -th term is 1 and of which all the other terms are 0. Show that there exists an integer $B > 1$ such that for any product F of (positive) Fibonacci numbers, there exists a finite sequence $s = (s_1, \dots, s_n)$ with all $s_i \in \{1, 2, \dots, B\}$ such that $p(s) = F$.

Problem B (folklore)

Let ℓ be a prime number. For any group homomorphism $f: A \rightarrow B$ between abelian groups and for any integer $n \geq 0$, denote by f_n the induced homomorphism $A/\ell^n A \rightarrow B/\ell^n B$. Let $(k_n)_{n=0}^\infty$ and $(c_n)_{n=0}^\infty$ be sequences of integers.

Show that there exist integers $N, a, b \geq 0$ and a group homomorphism $f: (\mathbb{Z}/\ell^N \mathbb{Z})^a \rightarrow (\mathbb{Z}/\ell^N \mathbb{Z})^b$ such that for all $n \geq 0$ we have $\#\ker f_n = \ell^{k_n}$ and $\#\operatorname{coker} f_n = \ell^{c_n}$ if and only if $k_0 = c_0 = 0$ and the sequences $(k_{n+1} - k_n)_{n=0}^\infty$ and $(c_{n+1} - c_n)_{n=0}^\infty$ are non-negative, non-increasing, eventually zero, and there is a constant C such that for all n such that $k_{n+1} - k_n$ and $c_{n+1} - c_n$ are not both zero, their difference is C .

(Recall that the *cokernel* $\operatorname{coker} f$ of a group homomorphism $f: A \rightarrow B$ between abelian groups is the quotient of B by the image of f .)

Problem C (folklore)

Let R be the polynomial ring over \mathbb{Z} with variables x_i, y_i, z_i for all $i \in \mathbb{Z}$. Let S be the polynomial ring over \mathbb{Z} with variables t_i for all $i \in \mathbb{Z}$. Let $\tau: R \rightarrow S$ be the isomorphism of rings given by $x_i \mapsto x_{i+1}, y_i \mapsto y_{i+1}$ and $z_i \mapsto z_{i+1}$.

Consider the morphism $f: R \rightarrow S$ of rings given by $x_i \mapsto t_{i-1} t_i t_{i+1}, y_i \mapsto t_i^3$ and $z_i \mapsto t_i^2$. Does there exist a finite number of elements $r_1, \dots, r_n \in R$ such that the kernel I of f is generated as an ideal in R by $\{\tau^i r_j; i \in \mathbb{Z}, j = 1, \dots, n\}$?

Edition 2016-2 We received solutions from Hendrik Reuvers, Pieter de Groen en Toshihiro Shimizu.

Problem 2016-2/A (folklore)

Denote for all positive rational numbers x by $f(x)$ the minimum number of 1's needed in a formula for x involving only ones, addition, subtraction, multiplication, division and parentheses. For example, $f(1) = 1$, and $f(\frac{1}{3}) = 4$, as $\frac{1}{3} = \frac{1}{1+1+1}$ and as no such formula exists with at most three 1's. Note that $f(11) \neq 2$ (concatenation of ones is not allowed). Moreover, denote for all positive rational numbers x by $h_2(x)$ the number $\log_2(p) + \log_2(q)$, where \log_2 denotes the base-2 logarithm, and where p, q are positive integers such that $x = \frac{p}{q}$ and $\gcd(p, q) = 1$.

Show that for all x , we have

$$f(x) > \frac{1}{2} h_2(x).$$

Redactie:

Gabriele Dalla Torre

Christophe Debry

Jinbi Jin

Marco Streng

Wouter Zomervrucht

Problemenrubriek NAW

Mathematisch Instituut

Universiteit Leiden

Postbus 9512

2300 RA Leiden

problems@nieuwarchief.nl

www.nieuwarchief.nl/problems

Oplösungen

| Solutions

Solution We received a solution from Toshihiro Shimizu, to whom the book token is awarded, and the solution of whom the following is based on.

Let x be a positive rational number, and let p, q be positive integers such that $x = \frac{p}{q}$ and $\gcd(p, q) = 1$. We show by induction on $f(x)$ that $p, q \leq 2^{f(x)-1}$; from this it will follow that $2f(x) > 2f(x) - 2 \geq \log_2 p + \log_2 q = h_2(x)$, as desired.

First note that $x = 1$ is the only rational number with $f(x) = 1$, and in this case $p = q = 1 = 2^{f(x)-1}$. So suppose that n is a positive integer and that for any x with $f(x) \leq n$ written as $\frac{p}{q}$ with p, q positive integers with $\gcd(p, q) = 1$, we have $p, q \leq 2^{f(x)-1}$. Suppose that $f(x) = n + 1$. Then x can be written in one of the forms $x_1 + x_2, x_1 - x_2, x_1 \cdot x_2, \frac{x_1}{x_2}$ with x_1, x_2 positive rational numbers with $f(x_1) + f(x_2) = f(x) = n + 1$ (so that $f(x_1), f(x_2) \leq n$). Write $x_i = \frac{p_i}{q_i}$ where p_i and q_i are positive integers such that $\gcd(p_i, q_i) = 1$, for $i = 1, 2$.

If $x = x_1 + x_2$ or $x = x_1 - x_2$, then we find that $x = \frac{p_1 q_2 \pm p_2 q_1}{q_1 q_2}$, so $p, q \leq 2^{1+f(x_1)-1+f(x_2)-1} = 2^n = 2^{f(x)-1}$. If $x = x_1 \cdot x_2$, then we find that $x = \frac{p_1 p_2}{q_1 q_2}$, so $p, q \leq 2^{f(x_1)-1+f(x_2)-1} = 2^{n-1} < 2^{f(x)-1}$.

The same argument shows that if $x = \frac{x_1}{x_2}$, that then $p, q < 2^{f(x)-1}$. Hence we always have $p, q \leq 2^{f(x)-1}$, and we are done.

Problem 2016-2/B (folklore)

Suppose that there are $N \geq 2$ players, labeled $1, 2, \dots, N$, and that each of them holds precisely $m \geq 1$ coins of value 1, m coins of (integer) value $n \geq 2$, m coins of value n^2 , et cetera. A *transaction* from player i to player j consists of player i giving a finite number of his coins to player j . We say that an N -tuple (a_1, a_2, \dots, a_N) of integers is (m, n) -payable if $\sum_{i=1}^N a_i = 0$ and after a finite number of transactions, the i -th player has received (in value) a_i more than he has given away.

Show that for every N -tuple (a_1, a_2, \dots, a_N) with $\sum_{i=1}^N a_i = 0$ to be (m, n) -payable, it is necessary and sufficient that $m > n - \frac{n}{N} - 1$.

Solution We received solutions from Pieter de Groen and Toshihiro Shimizu. The book token goes to Pieter de Groen. Both solutions shared the same idea, the following solution is based on that of Toshihiro Shimizu.

First observe that if an N -tuple (a_1, a_2, \dots, a_n) is (m, n) -payable, the number of coins of value 1 that player i receives is a_i modulo n , since coins of higher value are of value divisible by n . Also observe that $m > n - \frac{n}{N} - 1$ is equivalent to $mN \geq (n-1)(N-1)$.

Note that if $m \leq n - \frac{n}{N} - 1$ (so $m < n$), or equivalently, $mN < (n-1)(N-1)$, then the tuple

$$(n - m - 1, n - m - 1, \dots, n - m - 1, -(N - 1)(n - m - 1))$$

is not (m, n) -payable; every player up to player $N - 1$ has to either receive at least $n - m - 1$ coins of value 1, or give away at least $m + 1$ coins of value 1. The latter is clearly impossible. This means that the last player must give away at least $(N - 1)(n - m - 1)$ coins of value 1. However, $(N - 1)(n - m - 1) = (n - 1)(N - 1) - m(N - 1) > m$, so this is also impossible. Therefore the condition $m > n - \frac{n}{N} - 1$ is necessary.

Next, we show that the condition $m > n - \frac{n}{N} - 1$ is sufficient. Let (a_1, \dots, a_N) be any N -tuple of integers with $\sum_i a_i = 0$. Assume that $m > n - \frac{n}{N} - 1$. We show that (a_1, \dots, a_N) is (m, n) -payable, by induction on $\max_i |a_i|$.

If $\max_i |a_i| \leq 1$, then all a_i lie in $\{-1, 0, 1\}$, and as the sum of the a_i 's is zero, the tuple is (m, n) -payable (by having every player with negative a_i pay one coin of value 1, and every player with positive a_i receive one coin of value 1). So assume that $A > 1$, and that all tuples (a_1, \dots, a_N) with zero sum and $\max_i |a_i| < A$ are (m, n) -payable.

Let (a_1, \dots, a_N) be a tuple with $\max_i |a_i| = A$. Let r_i denote the remainder of a_i on division by n . We assume without loss of generality that $n > r_1 \geq r_2 \geq \dots \geq r_N \geq 0$. Note that $\sum_i r_i$ is a multiple of n , say $\sum_i r_i = kn$. We show that $n - r_1, \dots, n - r_k \leq m$, so that player i can pay $n - r_i$ coins of value 1 for $i = 1, 2, \dots, k$.

Note that

$$kn = \sum_i r_i = (r_1 + \dots + r_{k-1}) + (r_k + \dots + r_N) \leq (N - k + 1)r_k + (k - 1)n.$$

Therefore $r_k \geq \frac{n}{N - k + 1} \geq \frac{n}{N}$, so $n - r_k = n - \frac{n}{N} < m + 1$, from which we deduce that $n - r_k \leq m$, and therefore also that $n - r_i \leq m$ for all $i = 1, 2, \dots, k$. So by having, for $i = 1, 2, \dots, k$,

player i pay $n - r_i$, and for all other i , player i receive r_i (which is possible by the above and since $\sum_i r_i = kn$), we see that the tuple (a_1, \dots, a_N) is (m, n) -payable if $(a_1 + n - r_1, \dots, a_k + n - r_k, a_{k+1} - r_{k+1}, \dots, a_N - r_N)$ is (m, n) -payable using only coins of value n or higher, and therefore if the tuple

$$(a'_1, \dots, a'_N) = \left(\frac{1}{n}(a_1 + n - r_1), \dots, \frac{1}{n}(a_k + n - r_k), \frac{1}{n}(a_{k+1} - r_{k+1}), \dots, \frac{1}{n}(a_N - r_N)\right)$$

is (m, n) -payable.

We finish the induction by showing that $\max_i |a'_i| < A$, so that by the induction hypothesis, the tuple (a'_1, \dots, a'_N) is indeed (m, n) -payable. Note that $a'_i = \lfloor \frac{a_i}{n} \rfloor$ or $a'_i = \lceil \frac{a_i}{n} \rceil$, so $|a'_i| \leq \lceil \frac{|a_i|}{n} \rceil$. Since $\lceil \frac{|a_i|}{n} \rceil \leq |a_i|$ with equality if and only if $|a_i| = 0$ or $|a_i| = 1$, it follows that $|a'_i| < |a_i|$ for all i with $|a_i| \geq 2$. Since $A \geq 2$, it follows that $\max_i |a'_i| < A$, and we are done.

Problem 2016-2/C (proposed by Wouter Zomervrucht)

For each integer $n \geq 1$ let c_n be the largest real number such that for any finite set of vectors $X \subset \mathbb{R}^n$ with $\sum_{v \in X} |v| \geq 1$ there exists a subset $Y \subseteq X$ with $|\sum_{v \in Y} v| \geq c_n$. Prove the recurrence relation

$$c_1 = \frac{1}{2}, \quad c_{n+1} = \frac{1}{2\pi n c_n}.$$

Solution We received solutions from Hendrik Reuvers and Toshihiro Shimizu. The book token is awarded to Hendrik Reuvers. Both solutions shared the same idea, the following solution is based on the one sent in by the proposer.

For $n \geq 0$, let $D^n \subset \mathbb{R}^n$ be the closed unit ball and $S^n \subset \mathbb{R}^{n+1}$ the unit sphere. Denote by v_n the volume of D^n and by s_n the surface area of S^n . We will show that $c_n = v_{n-1}/s_{n-1}$, then we are done by the relations $v_0 = 1$, $s_0 = 2$, $v_n = \frac{1}{n}s_{n-1}$, and $s_n = 2\pi v_{n-1}$.

First we make a computation. Let $n \geq 1$ and write $V_+ = \{x \in \mathbb{R}^n : x_n \geq 0\}$. For $r > 0$ we let rD^n be the closed radius r ball, then one has

$$\int_{rD^n \cap V_+} x_n dx = \int_0^r x_n (r^2 - x_n^2)^{\frac{n-1}{2}} v_{n-1} dx_n = \frac{v_{n-1}}{n+1} r^{n+1},$$

so

$$\int_{S^{n-1} \cap V_+} x_n dx = \frac{d}{dr} \left[\int_{rD^n \cap V_+} x_n dx \right]_{r=1} = v_{n-1}.$$

Now we turn to the problem. Take $n \geq 1$ and any collection $X \subset \mathbb{R}^n$ with $\sum_{v \in X} |v| \geq 1$. Let $Y \subseteq X$ be a subset for which the subsum $w = \sum_{v \in Y} v$ has maximal norm. Then Y must contain all $v \in X$ with $v \cdot w > 0$, and no $v \in X$ with $v \cdot w < 0$. In fact, $|w| = m(w/|w|)$, where for any $x \in S^{n-1}$ we define

$$m(x) = \sum_{v \in X \text{ with } v \cdot x \geq 0} v \cdot x.$$

It follows that $|w| = \max_{x \in S^{n-1}} m(x)$. For $v \in \mathbb{R}^n$ set $V_v = \{x \in \mathbb{R}^n : v \cdot x \geq 0\}$. (For instance, $V_+ = V_{(0, \dots, 0, 1)}$.) We compute

$$\int_{S^{n-1}} m(x) dx = \sum_{v \in X} \int_{S^{n-1} \cap V_v} v \cdot x dx = \sum_{v \in X} |v| \int_{S^{n-1} \cap V_v} x_n dx \geq v_{n-1}.$$

Thus there is $x \in S^{n-1}$ where $m(x) \geq v_{n-1}/s_{n-1}$, hence $c_n \geq v_{n-1}/s_{n-1}$.

Conversely, let X consist of k vectors with lengths $1/k$ and directions distributed homogeneously over S^{n-1} . (There are several ways of doing this; thanks to Toshihiro Shimizu for pointing out [1].) As $k \rightarrow \infty$, the associated function m converges uniformly to a constant function; by the computations above, its constant value is v_{n-1}/s_{n-1} . So also $c_n \leq v_{n-1}/s_{n-1}$.

Reference

¹ Eric W. Weisstein, Sphere Point Picking, <http://mathworld.wolfram.com/SpherePointPicking.html>.